

## CONTINUING MEDICAL EDUCATION

### Clinical microbiological case: severe relapsing septal panniculitis in a healthy man from the south-eastern USA

A. Safdar<sup>1</sup>, P. L. McEvoy<sup>2</sup>, R. G. Burns<sup>3</sup> and J. R. Perfect<sup>4</sup>

<sup>1</sup>Division of Infectious Diseases, Department of Medicine, University of South Carolina School of Medicine, Columbia, SC, <sup>2</sup>Department of Infectious and Parasitic Diseases, Armed Forces Institute of Pathology, Washington, DC, <sup>3</sup>Department of Pathology, University of South Carolina School of Medicine, Columbia, SC and <sup>4</sup>Department of Medicine, Duke University Medical Center, Durham, NC, USA


Accepted 18 June 2002

#### CASE REPORT

A 34-year-old, healthy caucasian man presented with relapsing, painful nodular lesions involving both the upper and lower extremities. His symptoms commenced eight weeks after burying two domestic cats that had died suddenly. He complained of profound fatigue, generalized myalgia, and joint pain, although he denied having cough, fever, and night sweats. An 8-cm arciform lesion on the anterior-medial aspect of the right upper arm had bright erythematous borders and a violaceous center. He was afebrile with small, non-tender, generalized lymphadenopathy. Ophthalmologic examination was normal. His white blood

cell count was 11 700 cells/ $\mu$ L and his platelet count was 388 000/ $\mu$ L. Serum aminotransferase and ACE levels, and erythrocyte sedimentation rate were normal. His symptoms worsened after two weeks of oral methylprednisolone and cefuroxime axetil (500 mg twice daily) treatment.

Histologic examination of the right arm lesion showed severe panniculitis with prominent necrosis of subcutaneous fat (Figure 1). Stains for acid-fast bacilli (Ziehl-Neelsen), fungi (GMS), and bacteria (Brown-Hopps, Brown-Brenn), and Warthin-Starry stain, were negative. Blood and tissue cultures were sterile. CT scan showed a normal liver, spleen, and intrathoracic and abdominal lymph nodes. A whole-body gallium scan was normal.



**Figure 1** Extensive fat necrosis and septal panniculitis is observed (H&E, magnification  $\times 100$ ). There is no evidence of inflammatory infiltrates including histiocytes around small intralobular venules (arrow) or arteriole (arrow head).

Corresponding author and reprint requests: Amar Safdar, M.D., Department of Infectious Diseases, Infection Control, and Employee Health, 402 The University of Texas MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston Texas 77030, USA  
Tel: +1 713 794 4455  
Fax: +1 713 745 6339  
E-mail: [asafdar@mdanderson.org](mailto:asafdar@mdanderson.org)

## QUESTIONS

1. What is your clinical diagnosis?
2. What are the common diseases associated with this infection?
3. What cutaneous manifestations are frequently seen in this setting?
4. What is the optimal treatment?